

NEW MEXICO OIL CONSERVATION COMMISSION

SANTA FE, NEW MEXICO

MISCELLANEOUS NOTICES

Submit this notice in triplicate to the Oil Conservation Commission or its proper agent before the work specified is to begin. A copy will be returned to the sender on which will be given the approval, with any modifications considered advisable, or the rejection by the Commission or agent, of the plan submitted. The plan as approved should be followed, and work should not begin until approval is obtained. See additional instructions in the Rules and Regulations of the Commission.

Indicate nature of notice by checking below:

NOTICE OF INTENTION TO TEST CASING SHUT-OFF		NOTICE OF INTENTION TO SHOOT OR CHEMICALLY TREAT WELL	<input checked="" type="checkbox"/>
NOTICE OF INTENTION TO CHANGE PLANS		NOTICE OF INTENTION TO PULL OR OTHERWISE ALTER CASING	
NOTICE OF INTENTION TO REPAIR WELL		NOTICE OF INTENTION TO PLUG WELL	
NOTICE OF INTENTION TO DEEPEN WELL			

Artesia, New Mexico

Place

January 19th, 1949

Date

OIL CONSERVATION COMMISSION,
Santa Fe, New Mexico.

Gentlemen:

Following is a notice of intention to do certain work as described below at the

Robert E. McKee State Lease Well No. 10 in NE 1/4 NE 1/4 SW 1/4
Company of Operator
of Sec. 3, T. 19, R. 29, N. M. P. M., Turkey Track Field.
Eddy County.

FULL DETAILS OF PROPOSED PLAN OF WORK

FOLLOW INSTRUCTIONS IN THE RULES AND REGULATIONS OF THE COMMISSION

We intend to shoot this well in our lower pay zone between 2445 and 2489 feet. This zone will be shot with 200 quarts on January 20th, 1949.

Approved 1-27, 1949
except as follows:

Robert E. McKee

Company of Operator

By J. R. LumbPosition Authorized Agent

Send communications regarding well to

Name Robert E. McKeeAddress Box 246Artesia, New Mexico

OIL CONSERVATION COMMISSION,

By [Signature]Title Artesia Representative

1. The first part of the paper is devoted to the

study of the properties of the

operator T defined by the formula

$$Tf(x) = \int_0^x f(t) dt$$
 for $f \in L^p(\mathbb{R})$. It is shown that T is a bounded operator from $L^p(\mathbb{R})$ to $L^p(\mathbb{R})$ for $1 < p < \infty$. The norm of T is equal to 1. The operator T is also shown to be compact.

2. In the second part of the paper, the

operator T is extended to the space $L^p(\mathbb{R}^n)$ for $1 < p < \infty$. It is shown that T is a bounded operator from $L^p(\mathbb{R}^n)$ to $L^p(\mathbb{R}^n)$ for $1 < p < \infty$. The norm of T is equal to 1.

3. The third part of the paper is devoted to the

study of the properties of the operator T defined by the formula

$$Tf(x) = \int_0^x f(t) dt$$
 for $f \in L^p(\mathbb{R})$. It is shown that T is a bounded operator from $L^p(\mathbb{R})$ to $L^p(\mathbb{R})$ for $1 < p < \infty$. The norm of T is equal to 1.

4. The fourth part of the paper is devoted to the

study of the properties of the operator T defined by the formula

$$Tf(x) = \int_0^x f(t) dt$$
 for $f \in L^p(\mathbb{R})$. It is shown that T is a bounded operator from $L^p(\mathbb{R})$ to $L^p(\mathbb{R})$ for $1 < p < \infty$. The norm of T is equal to 1.

5. The fifth part of the paper is devoted to the

study of the properties of the operator T defined by the formula

$$Tf(x) = \int_0^x f(t) dt$$
 for $f \in L^p(\mathbb{R})$. It is shown that T is a bounded operator from $L^p(\mathbb{R})$ to $L^p(\mathbb{R})$ for $1 < p < \infty$. The norm of T is equal to 1.

6. The sixth part of the paper is devoted to the study of the properties of the operator T defined by the formula

$$Tf(x) = \int_0^x f(t) dt$$
 for $f \in L^p(\mathbb{R})$. It is shown that T is a bounded operator from $L^p(\mathbb{R})$ to $L^p(\mathbb{R})$ for $1 < p < \infty$. The norm of T is equal to 1.

7. The seventh part of the paper is devoted to the

study of the properties of the operator T defined by the formula

$$Tf(x) = \int_0^x f(t) dt$$
 for $f \in L^p(\mathbb{R})$. It is shown that T is a bounded operator from $L^p(\mathbb{R})$ to $L^p(\mathbb{R})$ for $1 < p < \infty$. The norm of T is equal to 1.

8. The eighth part of the paper is devoted to the

study of the properties of the operator T defined by the formula

$$Tf(x) = \int_0^x f(t) dt$$
 for $f \in L^p(\mathbb{R})$. It is shown that T is a bounded operator from $L^p(\mathbb{R})$ to $L^p(\mathbb{R})$ for $1 < p < \infty$. The norm of T is equal to 1.

9. The ninth part of the paper is devoted to the

study of the properties of the operator T defined by the formula

$$Tf(x) = \int_0^x f(t) dt$$
 for $f \in L^p(\mathbb{R})$. It is shown that T is a bounded operator from $L^p(\mathbb{R})$ to $L^p(\mathbb{R})$ for $1 < p < \infty$. The norm of T is equal to 1.

10. The tenth part of the paper is devoted to the

study of the properties of the operator T defined by the formula

$$Tf(x) = \int_0^x f(t) dt$$
 for $f \in L^p(\mathbb{R})$. It is shown that T is a bounded operator from $L^p(\mathbb{R})$ to $L^p(\mathbb{R})$ for $1 < p < \infty$. The norm of T is equal to 1.

11. The eleventh part of the paper is devoted to the

study of the properties of the operator T defined by the formula