

Address..... **Burt Bldg., Dallas Texas**

ACCEPTED

[illegible]

051111

1. The first group of authors (e.g., [1, 2]) considers the problem of the stability of the motion of a system of particles in the field of a central body. The results of the calculations show that the motion of the particles is stable in the case of a central body with a constant mass and a constant angular momentum. The results of the calculations show that the motion of the particles is stable in the case of a central body with a constant mass and a constant angular momentum.

Figure 1. Schematic representation of the experimental design. The subjects were divided into two groups: the control group and the experimental group. The control group was divided into two subgroups: the control group and the experimental group. The experimental group was divided into two subgroups: the control group and the experimental group. The control group was divided into two subgroups: the control group and the experimental group. The experimental group was divided into two subgroups: the control group and the experimental group.

[Faint handwritten notes and markings are visible at the bottom of the page.]

...and the fact that the *Journal* is a journal of the American Psychological Association, the largest and most influential organization in the field of psychology, adds to the journal's prestige and makes it a must-read for all psychologists.

Figure 1. Schematic representation of the experimental design. The subjects were divided into two groups: the control group and the experimental group. The control group was divided into two subgroups: the control group and the experimental group. The experimental group was divided into two subgroups: the control group and the experimental group. The control group was divided into two subgroups: the control group and the experimental group. The experimental group was divided into two subgroups: the control group and the experimental group.

Figure 1. A schematic diagram of the experimental setup. The subject is seated in a chair, viewing a video screen. The screen displays a target (a small circle) and a starting point (a small circle). The subject's hand is positioned at the starting point. The distance between the starting point and the target is 10 cm. The subject is instructed to move their hand from the starting point to the target. The video screen is positioned 40 cm from the subject's hand. The subject's hand is positioned at the starting point. The distance between the starting point and the target is 10 cm. The subject is instructed to move their hand from the starting point to the target. The video screen is positioned 40 cm from the subject's hand. The subject's hand is positioned at the starting point. The distance between the starting point and the target is 10 cm. The subject is instructed to move their hand from the starting point to the target. The video screen is positioned 40 cm from the subject's hand.

$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{x}} \right) = \frac{\partial L}{\partial x}$

Figure 1. The effect of the concentration of the initiator on the polymerization of α -methylstyrene in the presence of $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ and $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ at 50°C in CH_2Cl_2 . The concentration of $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ was 1.0×10^{-3} mol/L, and the concentration of $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ was 1.0×10^{-3} mol/L. The concentration of α -methylstyrene was 0.1 mol/L, and the concentration of the initiator was 0.001 mol/L. The reaction time was 24 h.