

ATTACHMENT TO FORM C-105
GALLEGOS CANYON UNIT WELL NO. 145

4-1/2" casing tested to 3500 psi, held ok. Perforated Dakota 6042-50 and 6058-66 with 2 shots per foot and 6076-80 with 4 shots per foot. Fracked with 38,200 gallons water containing 7 pounds J-114 per 1000 gallons and 1% calcium chloride and 36,500 pounds 20-40 mesh sand. There was no breakdown. Pressures were: Pumped in 700 psi; Maximum treating 3500 psi; Minimum treating 3400 psi; Average treating 3500 psi; Instant shut in 1700 psi; 5 minute shut in 1100 psi; stabilized shut in 225 psi in 2-1/2 hours. The average injection rate was 34 barrels per minute.

A bridge plug was set at 6005'. Perforated Graneros 5958-68 with 4 shots per foot. Fracked with 23,200 gallons water with 7 pounds J-114 per 1000 gallons and 1 pound calcium chloride and 15,000 pounds 20-40 mesh sand and 5,000 pounds 10-20 mesh sand. The average injection rate was 40 barrels per minute. Pressures were: Breakdown 2400 psi; Maximum treating 3300 psi; Minimum treating 3200 psi; Average treating 3200 psi; Instant shut in 1600 psi; 5 minute shut in 1100 psi; Stabilized shut in pressure 0 psi in 1-1/2 hours.

Tubing landed at 6030'. Potential test 6264 MCF per day. Completed as a shut in Basin Dakota Field Development gas well.



Injection rate was 10 barrels per minute.
and in 11 1/2 hrs. fully acidified and in 2 1/2 hrs. The average
and 3000 psi; Average treated 5000 psi; Injection rate in 2 1/2 hrs.
pressure as well. Pumped in 70 psi; maximum treated 1000 psi; Injection rate
calcium chloride and 30,000 pounds 10-10 mesh sand. There was no production
and 30,000 pounds water containing 2 pounds 1-1/2% per 1000 psi in 10
and 3000-60 with 2 shots per foot and 3000-80 with 4 shots per foot. Total
2-1/2 hrs. casing tested to 3500 psi, held 1000 psi. Perforated 200 ft. in 10-1/2 hrs.

1. The first group of patients (Group I) consisted of 10 patients who were treated with 100 mg of penicillin G (Keflin) intramuscularly (i.m.) four times a day for 10 days. The second group (Group II) consisted of 10 patients who were treated with 100 mg of penicillin G (Keflin) intramuscularly (i.m.) four times a day for 10 days. The third group (Group III) consisted of 10 patients who were treated with 100 mg of penicillin G (Keflin) intramuscularly (i.m.) four times a day for 10 days. The fourth group (Group IV) consisted of 10 patients who were treated with 100 mg of penicillin G (Keflin) intramuscularly (i.m.) four times a day for 10 days. The fifth group (Group V) consisted of 10 patients who were treated with 100 mg of penicillin G (Keflin) intramuscularly (i.m.) four times a day for 10 days. The sixth group (Group VI) consisted of 10 patients who were treated with 100 mg of penicillin G (Keflin) intramuscularly (i.m.) four times a day for 10 days. The seventh group (Group VII) consisted of 10 patients who were treated with 100 mg of penicillin G (Keflin) intramuscularly (i.m.) four times a day for 10 days. The eighth group (Group VIII) consisted of 10 patients who were treated with 100 mg of penicillin G (Keflin) intramuscularly (i.m.) four times a day for 10 days. The ninth group (Group IX) consisted of 10 patients who were treated with 100 mg of penicillin G (Keflin) intramuscularly (i.m.) four times a day for 10 days. The tenth group (Group X) consisted of 10 patients who were treated with 100 mg of penicillin G (Keflin) intramuscularly (i.m.) four times a day for 10 days.

interfered with my work. (Jed Friedman, 1980) In 1980, I was

