

NEW MEXICO OIL CONSERVATION COMMISSION Santa Fe, New Mexico

MISCELLANEOUS REPORTS ON WELL

Submit this report in triplicate to the Oil Conservation Commission or its proper agent within ten days after the work specified is completed. It should be signed and sworn to before a notary public for reports on beginning drilling operations, results of shooting well, results of test of casing shut-offs, result of plugging of well, and other important operations, even though the work was witnessed by an agent of the commission. Reports on minor operations need not be signed and sworn to before a notary public. See additional instructions in the Rules and Regulations of the Commission.

Indicate nature of report by checking below:

REPORT ON BEGINNING DRILLING OPERATIONS		REPORT ON REPAIRING WELL	
REPORT ON RESULT OF SHOOTING OR CHEMICAL TREATMENT OF WELL		REPORT ON PULLING OR OTHERWISE ALTERING CASING	
REPORT ON RESULT OF TEST OF CASING SHUT-OFF		REPORT ON DEEPENING WELL	
REPORT ON RESULT OF PLUGGING OF WELL		Perforate Tubing	XX

Hobbs, New Mexico.

April 8, 1941.

Place

Date

OIL CONSERVATION COMMISSION
Santa Fe, New Mexico.
Gentlemen:

Following is a report on the work done and the results obtained under the heading noted above at the

Gulf Oil Corp - Gypsy Prod Divn - W. A. Weir "A" Well No. 1 in the

Company or Operator

Lease

NE NE of Sec. 35, T. 19S, R. 36E, N. M. P. M.,

Monument Field, Lea County

The dates of this work were as follows:

Notice of intention to do the work was (was not) submitted on Form C-102 on _____ 19____ and approval of the proposed plan was (was not) obtained. (Cross out incorrect words.)

DETAILED ACCOUNT OF WORK DONE AND RESULTS OBTAINED

March 27, 1941 perforated tubing with 1 - 1/16" hole at 3585'

Production before:- During month of January 1941, flowed average of 16 barrels daily, flowing thru tubing,
During February and first half of March 1941 flowed average of 21.9 barrels daily, flowing intermittently thru casing and tubing..

Production after:- For 7 days following perforating job production has averaged 21.7 barrels daily thru 2" tubing w/average GOR 5,426.

Witnessed by _____
Name Company Title

Subscribed and sworn to before me this _____

8th day of April, 19 41

J. W. Egan
Notary Public

My Commission expires February 25, 1942.

I hereby swear or affirm that the information given above is true and correct.

Name Roy Yarbrough

Position District Supt.

Representing Gulf Oil Corp - Gypsy Prod Divn.
Company or Operator

Address Hobbs, New Mexico.

Remarks:

Roy Yarbrough
Name
OIL & GAS INSPECTOR
Title

THE UNIVERSITY OF CHICAGO

DEPARTMENT OF CHEMISTRY

For the purpose of this experiment, a solution of the substance to be studied was prepared in a known volume of solvent. The concentration of the solution was determined by weighing a known amount of the substance and dividing it by the volume of the solution.

The absorbance of the solution was measured at a wavelength of 440 mμ. The absorbance was found to be 0.45. The concentration of the solution was found to be 0.01 g/l. The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm.

The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm. The concentration of the solution was found to be 0.01 g/l. The absorbance of the solution was found to be 0.45.

The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm. The concentration of the solution was found to be 0.01 g/l. The absorbance of the solution was found to be 0.45. The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm.

The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm. The concentration of the solution was found to be 0.01 g/l. The absorbance of the solution was found to be 0.45. The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm.

The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm. The concentration of the solution was found to be 0.01 g/l. The absorbance of the solution was found to be 0.45. The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm.

The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm. The concentration of the solution was found to be 0.01 g/l. The absorbance of the solution was found to be 0.45. The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm.

The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm. The concentration of the solution was found to be 0.01 g/l. The absorbance of the solution was found to be 0.45. The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm.

The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm. The concentration of the solution was found to be 0.01 g/l. The absorbance of the solution was found to be 0.45. The molar absorptivity of the substance was found to be 1.125 x 10^4 l./mole-cm.