Additional

Information

Received: 12/21/2021

COG Operating LLC Hasta La Vista 8 Fee SWD #1 C-108 – Additional Attachment December 6, 2021

Statement Regarding Injection Interval and Confining Layers

The injection interval for the Hasta La Vista 8 Fee SWD 1 proposed well is within the Devonian and Silurian formations. This unit is made up of the Wristen Group and Fusselman Formation. The overall thickness of this interval is between 1,400 to 2,000 feet consisting of limestone and dolostone with significant primary and secondary porosity and permeability. Below the injection interval is the Montoya Formation, which is between 300 and 500 feet thick. Figure 1 shows the isopach map of the Wristen Group and Fusselman Formation along with known faults and proposed well location. Figure 2 shows the isopach map of the Montoya Formation.

Basement faults in the area are sourced from the Tectonic Map of Texas (Ewing, 1990). The Precambrian Basement Map (Frenzel et al, 1988) is the source of the Precambrian faults which is inferred from subsurface data. The proposed Hasta La Vista 8 Fee SWD 1 well is located 4.6 miles to the west of the nearest Precambrian fault and 0.8 miles to the east of the nearest Basement fault.

Overlying the Wristen Group is the Woodford Shale. The Woodford Shale in this area is between 200 to 300 feet thick, consisting of mostly shale and mudstone that act as a confining layer above the injection interval. Figure 3 shows the isopach map of the Woodford Formation, known faults and proposed well location. The underlying confining layer below the injection interval for the Hasta La Vista 8 Fee SWD 1 proposed well is the Simpson Group. The Simpson Group in this area is between 600 to 800 feet thick and consists of a series of shales, carbonates, and some sand. The Simpson Group interval acts as a confining layer below the injection interval due to the large thickness and the amount of shale sequences within this interval. Figure 4 shows the isopach map of the Simpson Group, known faults and proposed well location. The Woodford Shale and the significant amount of shale within the Simpson Group will likely prevent fluids from migrating outside of the injection interval in this well. There is an additional 400 to 600 feet of thickness of the Ellenburger Formation below the Simpson Group and above Precambrian Basement (shown in Figure 5).

Regards,

Carrie M. Martin Staff Geologist

ConocoPhillips

carrie.m.martin@conocophillips.com

Carrie Ullartin

432-221-0479

Fusselman/Wristen Group Isopach and Faults Map -103.6 -103.55 -103.65 225 34E 32E 33E 32.35 235 32E 33E 1600-Hasta La Vista 8 Fee SWD 1 32E 33E 34E 32.2 2000 250 32E 255 255 258 33E 34E 35E 32.1 32.1 265 265 265 32E 265 -103.4 -103.5 -103.45 -103.35 -103.3 -103.65 -103.6 -103.55 10miles LEGEND Surface name Silurian / Devonian Isopach Area Permian NM TOWNSHIPS Contour inc 200 Geographic datum GCS_North_American_1927 Precambrian Faults **Basement Faults** Projection NAD27 / Texas Central (US ft) O Proposed well surface location 1:200000

Figure 1: Silurian / Devonian (Fusselman Formation / Wristen Group) Isopach and Faults Map

Upper Ordovician (Montoya) Isopach and Faults Map -103.6 -103.55 -103.5 -103.45 -103.4 -103.35 32E 225 22S 35E 33E 34E 32.35 32.35 400 32.3 235 238 32E 235 33E 32.3 35E 32.25 vista 8 Fee SWD 1 Hasta La 245 32E 32.2 33E 34E 36E 400 32.15 255 255 32E 258 34E 35E 32.1 400. 32.05 26S 32E 265 265 -103.65 -103.6 -103.55 -103.5 -103.45 -103.4 -103.35 -103.3 10miles LEGEND Area Surface name Upper Ordovician (Montoya) Isopach Permian NM TOWNSHIPS Geographic datum GCS_North_American_1927 Contour inc - Precambrian Faults 100 **Basement Faults** Scale Projection NAD27 / Texas Central (USft) O Proposed well surface location 1:200000

Figure 2: Upper Ordovician (Montoya) Isopach and Faults Map




Figure 3: Late Devonian (Woodford) Isopach and Faults Map

Middle Ordovician (Simpson) Isopach and Faults Map -103.6 -103.55 -103.5 -103.65 -103.45 -103.4 -103.35 32E 225 225 33E 35E 32.35 32.35 32.3 235 235 32E 235 32.3 35E 800 32.25 Hasta La Vista 8 Fee SWI 245 245 32E 33E 32.2 35E 322 32.15 255 258 255 32E 255 33E 34E 35E 32.1 32. 26S 32E 265 26S 265 33E -103.65 -103.55 -103.5 -103.45 -103.4 -103.6 -103.35 -103.3 10miles LEGEND Area Surface name Middle Ordovician (Simpson) Isopach Permian NM TOWNSHIPS Geographic datum GCS_North_American_1927 Contour inc Precambrian Faults 200 **Basement Faults** Projection NAD27 / Texas Central (US ft) Scale Proposed well surface location 1:200000

Figure 4: Middle Ordovician (Simpson Group) Isopach and Faults Map

Lower Ordovician (Ellenburger) Isopach and Faults Map -103.4 -103.6 -103.45 -103.55 22S 34E 225 33E 35E 400-32,35 235 32E 235 33E 34E 35E Hasta La Vista 8 Fee SWD 32E 248 33E 32.2 34E 400 600 000 25S 32E 32.1 32.1 32.05 265 265 265 32E 265 -103.5 -103.45 -103.4 -103.35 -103.3 -103.55 -103.65 -103.6 10miles LEGEND Surface name Lower Ordovician (Ellenburger) Isopach Area Permian NM TOWNSHIPS Geographic datum GCS_North_American_1927 Contour inc Precambrian Faults 200 Basement Faults Projection NAD27 / Texas Central (US ft) Scale O Proposed well surface location 1:200000

Figure 5: Lower Ordovician (Ellenburger) Isopach and Faults Map

Statement Regarding Historical Review of Earthquakes in the Area

Historical review of earthquakes was conducted within a 50 km area around the Hasta La Vista 8 Fee SWD 1 proposed well. Seismic events were sourced from USGS earthquake catalog from January 1, 1970 to December 2, 2021 with a magnitude greater than 1.0. Table 1 shows all the seismic events within the 50 km area. There are a total 58 seismic events within this time-period ranging in magnitude from 1.0 to 4.6. Figure 6 shows the seismic event locations as red circles. The nearest earthquake to the Hasta La Vista 8 Fee SWD 1 proposed well is 4.12 miles to the west.

Regards,

Carrie M. Martin

Staff Geologist ConocoPhillips

carrie.m.martin@conocophillips.com

Carried Martin

432-221-0479

Table 1: Seismicity within 50 km of proposed SWD well (USGS source: Jan. 1970 – Dec. 2021)

Date	Origin Time GMT	Latitude	Longitude	Depth (km)	Magnitude
2021-12-01	8:56:17	31.9846	-103.8697	7.73	3
2021-11-13	4:53:34	32.4132	-103.5923	5.00	3.2
2021-10-25	5:08:04	32.2083	-103.0893	2.45	2
2021-10-19	18:49:23	31.9882	-103.8719	7.83	2.3
2021-10-19	11:45:27	31.9892	-103.8751	7.52	2
2021-10-14	22:05:11	31.9901	-103.8697	7.52	3.3
2021-10-13	21:02:19	32.2220	-103.0925	2.23	2.2
2021-09-22	0:25:44	32.0694	-103.7176	7.62	1.7
2021-09-21	8:15:44	32.0726	-103.7116	8.24	2.3
2021-09-21	4:19:43	32.0680	-103.7159	8.91	3.2
2021-09-20	23:31:46	32.2128	-103.0989	3.55	1.9
2021-09-20	17:46:29	32.2037	-103.0968	4.43	2.7
2021-09-14	21:52:41	32.1927	-103.0903	6.04	1.8
2021-09-10	11:51:14	32.0870	-103.9025	8.34	2.3
2021-09-10	11:20:39	32.1780	-103.1139	8.00	1.9
2021-09-10	2:44:26	32.1780	-103.1118	5.99	1.9
2021-09-09	22:04:39	32.1909	-103.0925	5.70	2.5
2021-09-09	21:47:19	31.9451	-103.3775	7.91	1.7
2021-09-09	14:47:54	31.9432	-103.3764	8.21	2.7

2021-09-09	10:51:06	31.9382	-103.3770	10.12	1.6
2021-09-07	8:29:09	32.2028	-103.0957	1.61	2.2
2021-09-05	11:59:29	32.1872	-103.1075	6.24	2
2021-09-03	8:47:54	32.2037	-103.0968	5.26	2
2021-09-01	17:52:40	32.2137	-103.0914	3.18	2.4
2021-09-01	8:54:10	32.2000	-103.0903	1.64	2
2021-09-01	7:19:23	32.2128	-103.0946	5.01	2.5
2021-08-31	10:17:19	32.2101	-103.1129	7.09	2.1
2021-08-17	22:24:07	32.0722	-103.7170	7.42	2
2021-08-15	14:56:22	32.0740	-103.7116	7.83	3.2
2021-08-04	6:53:57	32.0754	-103.7132	7.52	1.9
2021-08-04	3:06:21	32.0731	-103.7116	7.73	2.3
2021-07-31	11:29:49	32.0648	-103.7224	8.39	2.5
2021-07-28	23:50:22	32.0703	-103.7132	8.70	2.7
2021-07-28	19:20:39	32.0830	-103.7124	7.29	2.2
2021-07-19	11:23:25	32.0712	-103.7192	7.83	4
2021-07-15	8:13:04	32.0510	-103.7559	7.42	1.6
2021-07-01	19:49:12	31.9594	-103.7962	6.75	2.3
2021-07-01	2:51:28	32.0840	-103.7879	8.09	1.6
2021-06-25	10:04:25	32.0699	-103.7257	7.42	2
2021-06-16	16:10:53	31.9051	-103.4137	9.50	1.6
2021-06-08	6:34:48	31.9668	-103.7957	7.52	2.1
2021-05-10	4:36:40	32.0372	-103.7294	8.70	1.8
2021-04-28	14:56:38	32.0317	-103.7672	7.93	1.9
2021-04-04	13:02:13	31.9074	-103.4434	6.05	1.9
2021-03-19	14:18:19	32.0326	-103.7656	8.60	2.7
2021-03-19	8:02:15	32.0503	-103.7616	8.68	2.2
2021-03-19	2:43:27	32.0284	-103.7683	8.34	3.4
2020-11-08	21:19:55	31.8986	-103.4622	5.98	2.7
2020-07-14	4:44:04	32.0528	-103.7722	11.84	2.5
2020-04-23	4:34:53	32.0439	-103.7650	9.81	2.6
2019-12-30	3:01:14	32.0402	-103.7668	5.00	2.5
2019-11-28	12:52:42	32.0263	-103.7789	9.65	2.7
2019-10-21	11:58:57	32.2659	-103.4070	5.00	2.7
2017-05-03	17:47:21	32.0819	-103.0226	5.00	2.6
2012-03-18	10:57:22	32.2810	-103.8920	5.00	3.1
2001-06-02	1:55:54	32.3340	-103.1410	5.00	3.3
1992-01-02	11:45:36	32.3360	-103.1010	5.00	4.6
1984-12-04	20:36:36	32.2660	-103.5560	5.00	2.9

Historical Seismicity and Faults Map




Figure 5: Historical Seismicity (Jan. 1, 1970 to Dec. 2, 2021) and Faults Map