

NEW MEXICO OIL CONSERVATION COMMISSION
SANTA FE, NEW MEXICO

Form C-110
Revised 7/1/55

(File the original and 4 copies with the appropriate district office)

CERTIFICATE OF COMPLIANCE AND AUTHORIZATION
TO TRANSPORT OIL AND NATURAL GAS

Company or Operator Western Oil Fields, Inc. Lease Grier
Well No. 1 Unit Letter D S 31 T 16 R 31 Pool Square Lake
County Eddy Kind of Lease (State, Fed. or Patented) Federal
If well produces oil or condensate, give location of tanks: Unit S T R
Authorized Transporter of Oil or Condensate Continental Pipeline Company
Address Box 367, Artesia, New Mexico
(Give address to which approved copy of this form is to be sent)
Authorized Transporter of Gas No Gas
Address _____
(Give address to which approved copy of this form is to be sent)
If Gas is not being sold, give reasons and also explain its present disposition:
None produced.

Reasons for Filing: (Please check proper box) New Well ()
Change in Transporter of (Check One): Oil (x) Dry Gas () C'head () Condensate ()
Change in Ownership () Other ()
Remarks: _____
(Give explanation below)

The undersigned certifies that the Rules and Regulations of the Oil Conservation Commission have been complied with.

Executed this the _____ day of _____ 19 _____

By ORIGINAL
SIGNED BY R. H. SWARTHOUT

Approved _____ 19 _____

Title Petroleum Engineer

OIL CONSERVATION COMMISSION

Company Western Oil Fields, Inc.

By M. L. Armstrong

Address Box 1147

Title _____

Hobbs, New Mexico

OIL CONSERVATION COMMISSION
EXECUTIVE OFFICE

ADDITIONAL DISTRICT OFFICE

1. *Chlorophyll a* and *Chlorophyll b* were determined by the method of Lichtenthal and Whistler (1973). The total chlorophyll content was determined by the method of Arar and Cook (1977). The carotenoid content was determined by the method of Lichtenthal and Whistler (1973). The total carotenoid content was determined by the method of Arar and Cook (1977). The total protein content was determined by the method of Lowry (1956). The total lipid content was determined by the method of Bligh and Dyer (1959). The total carbohydrate content was determined by the method of Dubois and Gilles (1950). The total nucleic acid content was determined by the method of Burton (1956). The total ash content was determined by the method of AOAC (1970). The total moisture content was determined by the method of AOAC (1970). The total dry matter content was determined by the method of AOAC (1970). The total organic acid content was determined by the method of AOAC (1970). The total alkaloid content was determined by the method of AOAC (1970). The total saponin content was determined by the method of AOAC (1970). The total tannin content was determined by the method of AOAC (1970). The total flavonoid content was determined by the method of AOAC (1970). The total phenol content was determined by the method of AOAC (1970). The total terpenoid content was determined by the method of AOAC (1970). The total steroid content was determined by the method of AOAC (1970). The total glycoside content was determined by the method of AOAC (1970). The total alkaloid content was determined by the method of AOAC (1970). The total saponin content was determined by the method of AOAC (1970). The total tannin content was determined by the method of AOAC (1970). The total flavonoid content was determined by the method of AOAC (1970). The total phenol content was determined by the method of AOAC (1970). The total terpenoid content was determined by the method of AOAC (1970). The total steroid content was determined by the method of AOAC (1970). The total glycoside content was determined by the method of AOAC (1970).

File: