

OIL & GAS INSPECTOR

Page 15

live or neutral to site?

1. The following information is being furnished to you for your information and is not to be used for any other purpose. It is the property of the U.S. Government and is loaned to you. It is to be returned to the U.S. Government when requested to do so. It is to be kept confidential and not to be disclosed to any other person without the express written consent of the U.S. Government. It is to be destroyed when no longer needed for the purpose for which it was furnished. It is to be stored in a secure location and protected from unauthorized access. It is to be handled in accordance with the U.S. Government's security policies and procedures. It is to be used only for the purpose for which it was furnished. It is to be kept confidential and not to be disclosed to any other person without the express written consent of the U.S. Government. It is to be destroyed when no longer needed for the purpose for which it was furnished. It is to be stored in a secure location and protected from unauthorized access. It is to be handled in accordance with the U.S. Government's security policies and procedures. It is to be used only for the purpose for which it was furnished.

10. 11. 1950

Abstract. The purpose of this study was to determine whether there were differences in the prevalence of periodontitis between patients with type 2 diabetes mellitus (DM) and non-diabetic controls. A total of 60 patients with DM and 60 age- and sex-matched non-diabetic controls were recruited from a tertiary care hospital. All participants underwent a clinical examination of their periodontium by a single examiner. The prevalence of periodontitis was significantly higher among the diabetic group than among the control group ($P < .001$). The mean periodontal index score was significantly higher among the diabetic group than among the control group ($P < .001$). The results suggest that patients with DM have a higher prevalence of periodontitis compared with non-diabetic controls.

[illegible][illegible]

Abstract: The purpose of this study was to determine the effect of a 6-week training program on the physical fitness of female students. A total of 100 female students were divided into two groups: control and experimental. The control group did not receive any training, while the experimental group received a 6-week training program. Physical fitness was measured at baseline and after 6 weeks using a series of tests. The results showed that the experimental group had significantly higher scores than the control group in all tests after 6 weeks of training.

[illegible]

30-015-03803

WFX Order # 186