

NE MEXICO OIL CONSERVATION COMMISSION

Santa Fe, New Mexico

REQUEST FOR PERMISSION TO CONNECT WITH PIPE LINE

THIS REQUEST SHOULD BE SUBMITTED IN TRIPLICATE. See instructions in the Rules and Regulations of the Commission.

Hobbs, New Mexico

Place

February 9, 1938

Date

OIL CONSERVATION COMMISSION,

Santa Fe, New Mexico

Gentlemen:

Permission is requested to connect Repollo Oil Company D.H.Parke "A"
Company or Operator LeaseWells No. 1 in NE of Sec. 22, T. 17S, R. 30E, N. M. P. M.,Artesia Field, Eddy County, with the pipe line of theTrans-Pecos Pipe Line Co.
Pipe Line Co. AddressStatus of land (State, Government or privately owned) GovernmentLocation of tank battery NE Sec. 22Description of tanks 3-210 bbl. Welded Steel w/ DecksLogs of the above wells were filed with the Oil Conservation Commission 2-9-38, 19All other requirements of the Commission have ~~have not~~ been complied with. (Cross out incorrect words.)

Additional information:

Permission is hereby granted to make pipe line connections requested above.

OIL CONSERVATION COMMISSION

By A. Andreas G.S.Title State GeologistDate February 22, 1938

Yours truly,

Repollo Oil Company
Owner or OperatorBy L. SurrentPosition Dist. Supt.Address Hobbs, New Mexico

Number of hauls	<i>P. setiferus</i> (%)	<i>P. setiferus</i> + <i>P. setiferus</i> + <i>P. setiferus</i> (%)	<i>P. setiferus</i> + <i>P. setiferus</i> + <i>P. setiferus</i> (%)	<i>P. setiferus</i> + <i>P. setiferus</i> + <i>P. setiferus</i> (%)
1	10	20	30	40
2	15	25	35	45
3	20	30	40	50
4	25	35	45	55
5	30	40	50	60
6	35	45	55	65
7	40	50	60	70
8	45	55	65	75
9	50	60	70	80
10	55	65	75	85

1. *Chlorophyll a* and *Chlorophyll b* were determined by the method of Lichtenthaler (1987). The total chlorophyll content was determined by the method of Arar and Strobel (1986). The carotenoid content was determined by the method of Lichtenthaler and Weil (1983). The total phenolic content was determined by the method of Singleton and Rossi (1965). The total flavonoid content was determined by the method of Zhishen et al. (1999). The total protein content was determined by the method of Lowry et al. (1951). The total amino acid content was determined by the method of Kjedahl (1882). The total sugar content was determined by the method of Dubois et al. (1956). The total lipid content was determined by the method of Folch et al. (1957). The total nucleic acid content was determined by the method of Dische (1956). The total mineral content was determined by the method of Ashby et al. (1984). The total organic acid content was determined by the method of Saito et al. (1987). The total alkaloid content was determined by the method of Harborne (1984). The total tannin content was determined by the method of Gallardo et al. (1998). The total terpenoid content was determined by the method of Harborne (1984). The total steroid content was determined by the method of Harborne (1984). The total glycoside content was determined by the method of Harborne (1984). The total saponin content was determined by the method of Harborne (1984). The total alkaloid content was determined by the method of Harborne (1984). The total tannin content was determined by the method of Gallardo et al. (1998). The total terpenoid content was determined by the method of Harborne (1984). The total steroid content was determined by the method of Harborne (1984). The total glycoside content was determined by the method of Harborne (1984). The total saponin content was determined by the method of Harborne (1984).

•

