



**CHRONIC BRONCHITIS**

Chronic bronchitis is a disease of the bronchi, characterized by a persistent cough, often with sputum, and by a narrowing of the bronchi. It is usually caused by long-continued exposure to irritants, such as dust, smoke, or cold air, and is often associated with cigarette smoking.

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal. The only effective treatment is the removal of the irritant, and the avoidance of exposure to cold air.

CHRONIC BRONCHITIS

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal.

CHRONIC BRONCHITIS

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal.

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal.

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal.

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal.

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal.

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal.

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal.

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal.

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal.

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal.

The disease is usually progressive, and may lead to emphysema and finally to cor pulmonale. The treatment is symptomatic, and consists in the removal of the irritant, the use of expectorants, and the avoidance of exposure to cold air.

The prognosis is usually poor, and the disease is often fatal.