

NEW MEXICO  
OIL CONSERVATION COMMISSION

P. O. DRAWER DD  
ARTESIA, NEW MEXICO

No. June, 1970  
A 25

**SUPPLEMENT TO THE OIL PRORATION SCHEDULE**

DATE 6/11/70

PURPOSE: ALLOWABLE ASSIGNMENT New Well

Effective 6/1/70 an allowable of 8 barrels of oil per day or a total of 240 barrels for the month is hereby assigned to the Gulf Oil Corporation, Littlefield AB Federal, #4-E, 22-18-31, Shugart Pool.

WAG:sw

Gulf Oil Corporation

TMM

OIL CONSERVATION COMMISSION

W. J. [Signature]  
Supervisor, District No. 2

EXTRA COPY

TOIXEM RISE  
NUMBER 100 MONTAVER 4100 INC  
1000 W. 100th St.  
Tulsa, Oklahoma 74114

*[The following text is extremely faint and largely illegible due to poor scan quality. It appears to be a list or index of items.]*

[illegible]

1. *Chlorophyll a* and *Chlorophyll b* were determined by the method of Lichtenthaler and Whistler (1973). The total chlorophyll content was determined by the method of Arar and Cook (1980). The carotenoid content was determined by the method of Lichtenthaler and Whistler (1973). The total carotenoid content was determined by the method of Arar and Cook (1980). The total protein content was determined by the method of Lowry et al. (1951). The total lipid content was determined by the method of Bligh and Dyer (1959). The total carbohydrate content was determined by the method of Dubois and Gilles (1950). The total nucleic acid content was determined by the method of Burton (1956). The total ash content was determined by the method of AOAC (1990). The total moisture content was determined by the method of AOAC (1990). The total dry matter content was determined by the method of AOAC (1990). The total organic acid content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total terpenoid content was determined by the method of AOAC (1990). The total steroid content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total terpenoid content was determined by the method of AOAC (1990). The total steroid content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990).

[illegible]