

NEW MEXICO OIL CONSERVATION COMMISSION
INITIAL POTENTIAL TEST-DATA SHEET


This form must be used for reporting all pitot tube tests made in the State. It is particularly important that it be used for reporting Initial Potential Tests in the San Juan Basin as prescribed by Order No. R-333 and by the New Mexico Oil Conservation Commission Manual of Tables and Procedure for Initial Potential (Pitot Tube) Tests.

POOL Wildcat FORMATION Mesaverde
COUNTY Rio Arriba DATE WELL TESTED 9/3/56
Operator Magnolia Petroleum Lease Clearville Well No. H-1
Unit
1/4 Section NE Letter _____ Sec. 11 Twp. 26N Rge. 3W
Casing: _____ "O.D. Set At _____ Tubing _____ "WT. _____ Set At _____
Pay Zone: From _____ to _____ Gas Gravity: Meas. _____ Est. _____
Tested Through: Casing _____ Tubing X
Test Nipple 1.995 I.D. Type of Gauge Used X
(Spring) (Monometer)

OBSERVED DATA

Shut In Pressure: Casing _____ Tubing: 1700 S.I. Period 9-Day
Time Well Opened: 12:02 p.m. Time Well Gauged: 3:02 p.m.
Impact Pressure 52.04
Volume (Table I). 5726.8 (a)
Multiplier for Pipe or Casing (Table II).995 (b)
Multiplier for Flowing Temp. (Table III). 54.9 1.0058 (c)
Multiplier for SP. Gravity (Table IV). (d)
Ave. Barometer Pressure at Wellhead (Table V).
Multiplier for Barometric Pressure (Table VI). DIST. 3. (e)
Initial Potential, Mcf/24 hrs. (a) x (b) x (c) x (d) x (e) = 5731

Witnessed by _____ Tested by: W. J. McConathy *W. J. McConathy*
Company: _____ Company: Geoelectric, Inc
Title: _____ Title: Agent



OIL CONSERVATION COMMISSION
AZTEC DISTRICT OFFICE
No. 1001 - 1002

No. Copies Received

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

Journal of Interpersonal Violence 26(10) 1978–1996
© The Author(s) 2011
Reprints and permissions: <http://www.sagepub.com/journalsPermissions.nav>

Figure 1. The effect of the concentration of the *Agrobacterium* suspension on the transformation efficiency of *Agrobacterium* strains. The *Agrobacterium* strains were grown in the YEA medium for 24 h at 28°C. The cell concentration of the strains was adjusted to 10⁸ cells/ml. The cell suspension was mixed with the plant tissue and the transformation efficiency was determined. The results were expressed as the mean ± SD of three independent experiments. The asterisks indicate the significant difference between the strains.

100

[illegible]

100

...the