

REVISEDNEW MEXICO OIL CONSERVATION COMMISSION
Santa Fe, New Mexico

REQUEST FOR (OIL) - (GAS) ALLOWABLE

New Well
Recompletion

This form shall be submitted by the operator before an initial allowable will be assigned to any completed Oil or Gas well. Form C-104 is to be submitted in QUADRUPLICATE to the same District Office to which Form C-101 was sent. The allowable will be assigned effective 7:00 A.M. on date of completion or recompletion, provided this form is filed during calendar month of completion or recompletion. The completion date shall be that date in the case of an oil well when oil is delivered into the stock tanks. Gas must be reported on 15.025 psia at 60° Fahrenheit.

Hobbs, New MexicoSeptember 9, 1955

(Place)

(Date)

WE ARE HEREBY REQUESTING AN ALLOWABLE FOR A WELL KNOWN AS:

Pubco Development, Inc. Dickey State, Well No. 1-C, in NE $\frac{1}{4}$ SW $\frac{1}{4}$,
(Company or Operator) (Lease)

C, Sec. 31, T. 15 S, R. 31 E, NMPM, Unnamed Pool
(Unit)

Chaves County. Date Spudded 8-12-55, Date Completed 9-3-55

Please indicate location:

	X		

Elevation 4277.6 ground Total Depth 3086 driller, P.B. 3016 Schlum.Top oil/gas pay 2982 Sch. Top of Prod. Form Dickey-QueenCasing Perforations: 2991-2998 Sch. perf. log. orDepth to Casing shoe of Prod. String Casing set at 3059Natural Prod. Test 24 BOPDbased on 14 bbls. Oil in 14 Hrs. 0 Mins.Test after sand-oil free 429 BOPDBased on 222 143 bbls. Oil in 8 Hrs. 30 Mins.

Gas Well Potential

Size choke in inches Open 2"Date first oil run to tanks or gas to Transmission system: 9-3-55Transporter taking Oil or Gas: Artesia Pipeline Company

Casing and Cementing Record

Size Feet Sax

<u>9 5/8</u>	<u>236</u>	<u>175</u>
<u>5 1/2</u>	<u>3058.25</u>	<u>119</u>
<u>2" FUE</u>	<u>3008</u>	

Remarks:

I hereby certify that the information given above is true and complete to the best of my knowledge.

Approved: 1955, 19 PURCO DEVELOPMENT, INC.
(Company or Operator)

OIL CONSERVATION COMMISSION

By: C. M. KuderTitle Engineer District 1By: H. E. Maxwell, Jr. H. E. Maxwell Jr
(Signature)Title Supt. Prod. Dept.

Send Communications regarding well to:

Name H. E. Maxwell, JrAddress 107 West Chaco, Artesia, New Mexico

10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----

1. *Chlorophyll a* (Chl *a*)

[illegible]

$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{x}} \right) = \frac{\partial L}{\partial x}$

1. *Chlorophyll a* and *Chlorophyll b* were determined by the method of Lichtenthaler and Whistler (1973). The total chlorophyll content was determined by the method of Arar and Cook (1980). The carotenoid content was determined by the method of Lichtenthaler and Whistler (1973). The total carotenoid content was determined by the method of Arar and Cook (1980). The total protein content was determined by the method of Lowry et al. (1951). The total lipid content was determined by the method of Bligh and Dyer (1959). The total carbohydrate content was determined by the method of Dubois and Gilles (1950). The total nucleic acid content was determined by the method of Burton (1956). The total ash content was determined by the method of AOAC (1990). The total moisture content was determined by the method of AOAC (1990). The total dry matter content was determined by the method of AOAC (1990). The total organic acid content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total terpenoid content was determined by the method of AOAC (1990). The total steroid content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total terpenoid content was determined by the method of AOAC (1990). The total steroid content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990).