

FORMATION RECORD

| FROM | TO | THICKNESS IN FEET | FORMATION |
|---------|-------|----------------------|--|
| Surface | 328 | 328 | Caliche and surface clay. |
| 328 | 2205 | 1877 | Red beds and streaks of sand |
| 2205 | 3030 | 825 | Red beds with streaks anhydrite & sand |
| 3030 | 4415 | 1385 | Red beds with streaks sand and salt. |
| 4415 | 7799 | 3384 | Dolomite with streaks of sand. |
| 7799 | 9086 | 1287 | Dolomite with streaks of shale. |
| 9086 | 9350 | 324 | Lime & dolomite with streaks of shale & chert. |
| 9350 | 11015 | 1665 | Lime with streaks shale, chert, & sand. |
| 11015 | 11750 | 735 | Lime & shale with streaks of chert. |
| 11750 | 11978 | 228 | Dolomite |
| | 11978 | | Total Depth (Schlumberger). |
| | 11985 | 7 | Total Depth (Driller). |

FORMATION TOPS FROM SCHLUMBERGER

| | | |
|-------------------|--------|-----------|
| Top Anhydrite | 2205' | (+ 1694') |
| Top Yates | 3030' | (+ 869') |
| Top San Andres | 4415' | (- 516') |
| Top Glorietta | 5860' | (- 1961') |
| Top Tubbs | 7103' | (- 3204') |
| Top Abo | 7799' | (- 3900') |
| Top Wolfcamp | 9086' | (- 5187') |
| Top Pennsylvanian | 9793' | (- 5894') |
| Top Mississippian | 11015' | (- 7116') |
| Top Devonian | 11750' | (- 7851') |