


SCOUT REPORT NEW MEXICO

OIL CONSERVATION COMMISSION



Company Continental
 Farm Name State K-29 Well No. 1
 Land Classification State
 Sec. 29 Twp. 16 Range 37 County Lea
 Feet from Line: N. 660 S. E. 660 W.
 Elevation _____ Method Rotary
 Contractor Herchbach Drilling Co. Scout _____
 Spudded 6-9-38 Completed 8-31-38 Initial Production _____
 Bond Status _____

ACID RECORD

Casing and Cementing Record

Size	Feet	Inches	Sax Cement
12	251		200
9 7/8	2260		425

Gals.

Prod.	Date

Top Pay

Total Depth

SHOOTING RECORD

No. of Quarts	Shot at	Feet

Date		Date	
6-14	T.D.C.	AUG 9 - 1938	T.D. 4977 L
JUN 21 1938	T.D. 2876 K & A fractured pipe		Coring
JUN 22 1938	3356 R & GYP.	AUG 16 1938	T.D. 45217 L
7-5	A 4200 A		Open 4497-96
JUL 12 1938	Q 4587 L & A		Rec 7'6"
JUL 19 1938	T.D. 4905 L		12 hrs 55 min
	Prep to DST		D.S.T. 4917-96
	Core 4870-95	AUG 23 1938	T.D. 5202 L
	Blkly Core 4899-99 1/2		Blanchard
JUL 26 1938	T.D. 4915 L		Total Depth
	Core Split Assy.		No Oil Flow
	No Oil Gas nH ₂ O		
	on DST		
AUG 2 - 1938	T.D. 4915 WOC		
	Reconnect 30 sacks		

1. *Chlorophyll a* and *Chlorophyll b* were determined by the method of Lichtenthaler and Whistler (1973). The total chlorophyll content was determined by the method of Arar and Cook (1980). The carotenoid content was determined by the method of Lichtenthaler and Whistler (1973). The total carotenoid content was determined by the method of Arar and Cook (1980). The total protein content was determined by the method of Lowry et al. (1951). The total lipid content was determined by the method of Bligh and Dyer (1959). The total carbohydrate content was determined by the method of Dubois and Gilles (1950). The total nucleic acid content was determined by the method of Burton (1956). The total ash content was determined by the method of AOAC (1990). The total moisture content was determined by the method of AOAC (1990). The total dry matter content was determined by the method of AOAC (1990). The total organic acid content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total sterol content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total sterol content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990).

[illegible]