

NEW MEXICO OIL CONSERVATION COMMISSION

Santa Fe, New Mexico

MISCELLANEOUS NOTICES

Submit this notice in triplicate to the Oil Conservation Commission or its proper agent before the work specified is to begin. A copy will be returned to the sender on which will be given the approval, with any modifications considered advisable, or the rejection by the Commission or agent, of the plan submitted. The plan as approved should be followed, and work should not begin until approval is obtained. See additional instructions in the Rules and Regulations of the Commission.

Indicate nature of notice by checking below:

NOTICE OF INTENTION TO TEST CASING SHUT-OFF	<input checked="" type="checkbox"/>	NOTICE OF INTENTION TO SHOOT OR CHEMICALLY TREAT WELL	
NOTICE OF INTENTION TO CHANGE PLANS		NOTICE OF INTENTION TO PULL OR OTHERWISE ALTER CASING	
NOTICE OF INTENTION TO REPAIR WELL			
NOTICE OF INTENTION TO DEEPEN WELL		NOTICE OF INTENTION TO PLUG WELL	

Hobbs, New MexicoOctober 6, 1939

Place

Date

OIL CONSERVATION COMMISSION,
Santa Fe, New Mexico.

Gentlemen:

Following is a notice of intention to do certain work as described below at the

Cities Service Oil CompanyState-KWell No. 3 in NWNE

Company or Operator


Lease

of Sec. 27, T. 17, R. 35, N. M. P. M., Vacuum Field,Lea County.

FULL DETAILS OF PROPOSED PLAN OF WORK

FOLLOW INSTRUCTIONS IN THE RULES AND REGULATIONS OF THE COMMISSION

5½" casing was set at 4192' and cemented with 150 sax on October 5,
1939. We will let cement set for 72 hours and test for casing shut off.



OCT 9 - 1939

Approved _____, 19____
except as follows:

Cities Service Oil Company
Company or Operator

By D. D. Boile

Position Division Clerk
Send communications regarding well to

Name D. D. BoileAddress Hobbs, New Mexico

OIL CONSERVATION COMMISSION

By Roy YarrowTitle OIL & GAS INSPECTOR

Reliability of the Test

The reliability of the test is a measure of the consistency of the results. It is the degree to which the test results are the same when the test is repeated under the same conditions.

There are two main types of reliability: internal consistency and test-retest reliability. Internal consistency is the degree to which the items on the test measure the same construct. Test-retest reliability is the degree to which the test results are the same when the test is repeated at a later date.

Internal consistency is measured by the Cronbach's alpha coefficient. This coefficient ranges from 0 to 1, with 1 indicating perfect internal consistency. A Cronbach's alpha of 0.7 or higher is generally considered acceptable for most tests.

Test-retest reliability is measured by the Pearson correlation coefficient. This coefficient ranges from -1 to 1, with 1 indicating perfect test-retest reliability. A Pearson correlation of 0.7 or higher is generally considered acceptable for most tests.

There are several factors that can affect the reliability of a test. These factors include the quality of the test items, the stability of the test environment, and the characteristics of the test takers.

One way to improve the reliability of a test is to use a larger number of items. This will increase the internal consistency of the test and, therefore, the Cronbach's alpha coefficient.

Another way to improve the reliability of a test is to use a more stable test environment. This will reduce the variability in the test results and, therefore, increase the test-retest reliability.

Finally, it is important to use a representative sample of test takers. This will ensure that the test results are generalizable to the population of interest.

Overall, the reliability of a test is a critical factor in determining its validity. By following the guidelines outlined above, you can ensure that your test is reliable and that the results are consistent and generalizable.

There are several ways to improve the reliability of a test. These ways include using a larger number of items, using a more stable test environment, and using a representative sample of test takers.

By following these guidelines, you can ensure that your test is reliable and that the results are consistent and generalizable. This will help you to make more accurate decisions based on the test results.