

OIL CONSERVATION COMMISSION

BOX 2045

HOBBS, NEW MEXICO

DATE February 6, 1956

MR. W. B. MACEY
OIL CONSERVATION COMMISSION
P. O. BOX 871
SANTA FE, NEW MEXICO

RE: PROPOSED ORDER NO. DC 275

Dear Mr. Macey:

I have examined the application for dual completion dated 1/24
for Gulf R. R. Bell G #2-0 13-20-36
Operator Lease Name Well No. Unit S-T-R

and my recommendations are as follows:

OK CR AND RFM

Yours very truly,

OIL CONSERVATION COMMISSION

Engineer-District 1

1. The first part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation

$$f(x) = \int_0^x \frac{1}{1+t^2} dt$$

It is well known that the function $f(x)$ is increasing and concave down on the interval $(-\infty, \infty)$.

Let us consider the function $f(x)$ on the interval $(-\infty, \infty)$. It is well known that the function $f(x)$ is increasing and concave down on the interval $(-\infty, \infty)$. Let us consider the function $f(x)$ on the interval $(-\infty, \infty)$. It is well known that the function $f(x)$ is increasing and concave down on the interval $(-\infty, \infty)$.

Let us consider the function $f(x)$ on the interval $(-\infty, \infty)$. It is well known that the function $f(x)$ is increasing and concave down on the interval $(-\infty, \infty)$.

Let us consider the function $f(x)$ on the interval $(-\infty, \infty)$. It is well known that the function $f(x)$ is increasing and concave down on the interval $(-\infty, \infty)$.

Let us consider the function $f(x)$ on the interval $(-\infty, \infty)$. It is well known that the function $f(x)$ is increasing and concave down on the interval $(-\infty, \infty)$.

Let us consider the function $f(x)$ on the interval $(-\infty, \infty)$. It is well known that the function $f(x)$ is increasing and concave down on the interval $(-\infty, \infty)$.

Let us consider the function $f(x)$ on the interval $(-\infty, \infty)$. It is well known that the function $f(x)$ is increasing and concave down on the interval $(-\infty, \infty)$.

Let us consider the function $f(x)$ on the interval $(-\infty, \infty)$. It is well known that the function $f(x)$ is increasing and concave down on the interval $(-\infty, \infty)$.

Let us consider the function $f(x)$ on the interval $(-\infty, \infty)$. It is well known that the function $f(x)$ is increasing and concave down on the interval $(-\infty, \infty)$.

Let us consider the function $f(x)$ on the interval $(-\infty, \infty)$. It is well known that the function $f(x)$ is increasing and concave down on the interval $(-\infty, \infty)$.

$$f(x) = \int_0^x \frac{1}{1+t^2} dt$$

$$f(x) = \int_0^x \frac{1}{1+t^2} dt$$

$$f(x) = \int_0^x \frac{1}{1+t^2} dt$$