

2020

Age Group	Total (%)	Male (%)	Female (%)	Unknown (%)
18-24	100	45	55	100
25-34	100	55	45	100
35-44	100	45	55	100
45-54	100	35	65	100
55-64	100	25	75	100
65+	100	15	85	100

5

South curice

*

x

results obtained

Author Budge Drilling Company spudded 12-1/4" hole 6:20 2-2-57.

ran 10 joints (403') 8-5/8" of 24# Gr J-55 30 casing. Set and cemented at 414' with 300 sacks regular Best Cement. Plug at 399'. Maximum Pressure 250#. Circulated approximately 25 sacks cement. Job completed 10:45 2-3-57.

After waiting over 24 hours, tested 8-5/8" casing with 1000# for 30 minutes. No drop in pressure. Drilled cement plug from 379-414'. Tested below casing shoe with 500# for 30 minutes. No drop in pressure.

FILED IN CASE NO. FD-14 JUL 27 1968

Original Document

[illegible]

1. *Chlorophyll a* and *b* contents were determined by the method of Arar and Collins (1971).

Open, free, and fair trade is the only way to ensure that

RUSSEL, B. C. - 1960-1978

[REDACTED]

1) $\frac{1}{2} \leq \alpha \leq 1$ and $\frac{1}{2} \leq \beta \leq 1$ and $\alpha + \beta \leq 1$

Q. How many times did you see the defendant in the last 12 months?

Figure 1. The effect of the initial concentration of the monomer on the polymerization of α -methylstyrene initiated by TiCl_4 in CH_2Cl_2 at -78°C . The polymerization was carried out in the presence of 0.01 mole of TiCl_4 and 0.01 mole of CH_2Cl_2 in 10 ml of CH_2Cl_2 . The initial concentration of the monomer was varied from 0.01 to 0.1 mole/l. The polymerization was carried out for 10 min. The polymerization was carried out in the presence of 0.01 mole of TiCl_4 and 0.01 mole of CH_2Cl_2 in 10 ml of CH_2Cl_2 . The initial concentration of the monomer was varied from 0.01 to 0.1 mole/l. The polymerization was carried out for 10 min.

[illegible]

Wittneben et al.

Journal of Management Inquiry 20(4) 409-424

[illegible]

Area Supt. of Prod.
Gulf Oil Corporation