

SUNDRY NOTICES AND REPORTS ON WELLS

COPIES

[illegible]

Tulsa Oil & Cattle Co.

▲ 1955 年 12 月 20 日

4402 W. Industrial, Midland, Texas 79703

Indicators of child development clearly add to assessment, with any State required to develop a system to monitor child development.

449' 191 5 1 20' 721, Section 20

N/A

3458' GL

Check appropriate Box To Indicate Nature of Notice, Report, or Other Data

• • • • •

1. THE UNITED STATES OF AMERICA

[illegible]

MODEL NUMBER	FRANKLIN 5000
FRUIT OR VEGETABLE	STRAWBERRY
FRUITING OF PLANT	ABSENT
DATE	5-2-51
NAME OF GROWER	W. J. ...
ADDRESS	...
CITY	...
COUNTY	...
STATE	...
REMARKS	...

1. Lot 1, -3/4" 554 rubberer casing @ 40' on 4-30-84 mt. to surface w/ 20 ex.

$$f_{\text{max}} = \frac{1}{2\pi} \sqrt{\frac{1}{L C_{\text{eff}}}} = \frac{1}{2\pi} \sqrt{\frac{1}{L (C_1 + C_2)}} \quad (1)$$

3.0 Spud hole 10 ft. dia. 0 in. drilled to 400', ran 8-1/2" log cal. to 400'. Int. wt. 8.1 gal. in this Class "C" - 21 ea gal (14.8 gal/gal, yield 1.1 in. 3' x 100' total = 114 ft. 3) Class rate 7" cm. to surface. Plug down 3 9:10 a.m. 3 8.54. m. 10. 10. 10.

1. *Chlorophyll a* (Chl *a*) and *Chlorophyll b* (Chl *b*) were determined by the method of Lichtenthaler and Whistler (1973). The total chlorophyll content was determined by the method of Arar and Cook (1980). The carotenoid content was determined by the method of Lichtenthaler and Whistler (1973). The total carotenoid content was determined by the method of Arar and Cook (1980). The total protein content was determined by the method of Lowry (1956). The total lipid content was determined by the method of Bligh and Dyer (1959). The total carbohydrate content was determined by the method of Dubois and Gilles (1950). The total nucleic acid content was determined by the method of Burton (1956). The total ash content was determined by the method of AOAC (1990). The total dry weight was determined by the method of AOAC (1990). The total water content was determined by the method of AOAC (1990). The total organic acid content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total sterol content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total sterol content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990).

1. *Journal of the American Statistical Association*, 66, 1971, 107-114.

As a result, the model is able to capture the nonlinear relationship between the variables and the response variable, and it is able to handle the non-normal distribution of the response variable. The model is able to capture the nonlinear relationship between the variables and the response variable, and it is able to handle the non-normal distribution of the response variable.

Wm. A. Brown

ACCEPTED FOR RECORD

DATE OF: 17
SIGNATURES OF APPROVAL IF ANY:

JUN 18 1984

NEW MEXICO

*See Instructions on Reverse Side

1
RECEIVED
JUN 19 1984
F.C.E.
HIGHER OFFICE