

DESERT PAVEMENT

Not shown on map. Consists of a thin layer of closely spaced stones or rounded, over a surface layer of sand and silt. Stones collect at the surface by a sorting action, apparently due to wind and salt heaving, or swirling and shrinking of clay. It is beneath the pavement may be partly eolian in origin. In general, within a given area the pavement may be partly eolian in origin. In general, within a given area the pavement may be partly eolian in origin. In general, within a given area the pavement may be partly eolian in origin.

CAVE DEPOSITS

Not shown on map. Commonly have a well at base, recording an early stage of substantial water flow that eroded the cave. The gravel is overlain by clay or silt deposited as the flow of water diminished, and this in turn is overlain by stalagmites. Stalagmites are formed by dust. Fossil remains of Pleistocene animals may occur in deposits below the stalagmites. Remains of Holocene animals characterize the overlying deposits. Other cave deposits occur in basaltic lava, especially in the area southwest of the Zuni Mountains. These deposits include blocks fallen from the roofs, dust, and some ice.

ORGANIC DEPOSITS

Not shown on map. Accumulation of fibrous peat in ridge marshes bordering many flow areas. Both fibrous and woody peat accumulated in small, poorly drained depressions and more than 10 inches. Mostly less than 15 ft.

DESERT VARNISH

Not shown on map. A black stain of iron and manganese oxides on bare rock surfaces and on pebbles of desert pavement. Preserves prehistoric pottery-bearing occupations of the region. Predominantly middle Holocene, partly late Pleistocene. Many of these stained surfaces have petroglyphs carved by prehistoric peoples.

TRANSITIONAL DEPOSITS

Deposits transitional between those formed in situ and those transported; deposits moved downslope chiefly by gravity, particularly slow creep (colluvium). Also includes rock falls, landslides and avalanches shown as periglacial features.

Colluvium includes the heterogeneous mantle of soil and rock fragments derived from weathering of bedrock, and/or unconsolidated surficial deposits moved slowly downslope by gravitational force and sheet wash. Slopes generally steeper than 20 percent. Mass wasting, the process of water-saturated debris, frost heaving, alternate wetting and drying of clays, crystallization of salts, growth of roots, burrowing and trampling by animals, falling of trees, and impact of hail or rain. These, like other erosional processes, may be accelerated by man's activities.

Colluvium is basically a chaotic mixture of angular rock fragments and finer grained material. In New Mexico colluvium is generally less than 10 ft thick (locally 25 ft or more) but may grade into thick cones of debris at bases of hillsides. In the northeast and northwest parts of the state where steep shale slopes underlie the great caprock of sandstone or lava, two, and locally three, ages of colluvium may be distinguished. These are thought to be mid-Holocene, late Wisconsinan, and early Wisconsinan, respectively. Such occurrences provide an index of retreat of cliffs. Some shale slopes are armored and protected against erosion by blocks of the caprock.

On long low slopes such as flanks of the Zuni Mountains and east flank of the Sacramento Mountains, the colluvium is generally thin (commonly 1 to 2 ft thick) except near the bases of steep hillsides, and is composed of the resistant rock, forming the dip slope. Some of this colluvium could as well be mapped as rocky residue over limestone. Hillsides on granitic and volcanic rocks may also be overlain by thin but bouldery sandy colluvium. Colluvium on steep, faulted mountain fronts consists of a mixture of stones representing all the exposed formations up slope.

coll COLLUVIUM — Subscripts indicate the underlying hillside formation (e.g., collv. colluvium on Tertiary volcanic rocks)

TRANSPORTED DEPOSITS

Most surficial deposits are rocks and particles weathered from bedrock in one area, transported by water, wind, ice, or gravity to an area of deposition, and are susceptible to further erosion and transportation. These deposits are much younger than, and unrelated to, the underlying bedrock. They are classified according to their mode of transportation to the site of deposition.

ALLUVIUM IN FLOODPLAINS AND STREAM CHANNELS

Well stratified, sandy and silty stream deposits with gravel lenses; gravel lenses along valley walls. Consist of alluvial deposits record complex response to climatic change, shifts in local Mexican climate were comparatively wet during the Pleistocene glacial stage. Conversely, during the interglaciations, climates were drier, with conditions similar to Holocene environments. Alluvial deposits locally contain fossils, including bones of mammals and rodents, and shells of freshwater snails and clams. Late Pleistocene deposits contain fossil remains of extinct animals such as elephants, camels, horses, first reintroduced until the arrival of the Spaniards), sloths, and long-haired horses. Archaeological remains are common and on Holocene deposits and help date them. Three ages of alluvium generally can be distinguished: late Pleistocene, mid-Holocene, and historic. At least three recognized types of alluvial floodplain deposits reflect relative complexity of sediment transport by the main stream and its tributaries. A fourth type, along the Pecos River in the southwestern part of the state, is characterized by coarse ground, a fifth it is restricted to basalt-capped mesas.

al FLOODPLAIN AND CHANNEL DEPOSITS ALONG MAIN STREAMS — Generally very flat but includes terraces to about 10 ft high, shallow curved swales at right meanders, and local stabilized dunes. V. silty sand, silt, and some layers of gravel. Caliche absent or weakly developed in thin varieties; fibers, coatings on rocks, and soft nodules. Deposits commonly 25 ft thick. Ground water shallow, subject to pollution. Extensively farmed; subject to flooding.

al₂ FLOODPLAIN AND CHANNEL DEPOSITS ALONG GENERALLY PERMANENT RIVERS AND WASHES — Includes deposits along some permanent rivers and washes. Exposed to erosion and to erosion. Surface flat where deposit was formed by stream overflowing its banks; hummocky where sand of eroding fans at mouths of tributaries that crowd the main stream deposit at far bank. V. silty. Caliche common alluvium grades laterally into fan sand washed from eroding hillside. Lenticular perched water tables under some deposits. 15 ft. of deposits a possible has been exaggerated but total area probably about right because much of it is not to be omitted.

al₃ COARSE SILEY AND SANDY ALLUVIAL FANS — Intermediate between al₁ and some of the deposits (s and si)

al₄ SALTY ALLUVIUM — Borders Pecos River south of Fort Sumner

al/b ALLUVIUM OVER BASALT — Restricted to hillcapped mesas, strong, common, alluvium in old valleys, thickness commonly 10 ft or more. Acid soil



Gravel Terraces — Well-rounded stream gravels with cobbles 6 inches or more in diameter; some terraces 250 ft higher than the streams. Especially well developed along the San Juan River, less so along the Pecos, Gila, and Canadian Rivers. Most represent deposits by Pleistocene melt caliche deposits, especially on the higher terraces, which may be Kansan; lowest are Wisconsinan

ALLUVIAL FAN DEPOSITS

In alluvial fans, unlike floodplain alluvium, beds tend to be thick, massive, and highly lenticular rather than well stratified. This is characteristic of all the facies, whether boulder, gravel, sand, or silt. Beds lenticular and elongated down the slope of the fans; slopes 2 to 20 percent. Deposition mostly by flash floods, with poor sorting and mixed textures. Coarse-textured lenses commonly form ridges extending down the fan onto generally finer grained sediment. Boundaries between the textural facies of the deposits roughly parallel the fan contour, but detailed boundaries are irregularly lobate; those shown are approximations. Fan textures and slopes depend partly on composition of the parent rocks and partly on height and steepness of the bordering hill or mountain. Fans extensive in the Basin and Range part of the state where they comprise about half the total area; in other parts of the state, fans are small. On the larger fans, arroyos become shallower towards the toe; many head at low mounds that probably mark old mudflows. Ground subject to sheet flooding



Gravel Facies — Bouldery towards apex of fan, grading downslope to cobble and fine gravel with increasing proportion of sand and finer grained material. Commonly dissected to form 2 to 3 levels of gravel benches up to 50 ft above present washes. A few streams (e.g., Mulligan Wash, Alamogordo River, Cuchillo Negro Creek, and Rincon Arroyo) are incised 100 ft below fan surfaces. On short, steep fans, depths of valleys generally decrease downslope. On the broad Palomas surface, west of the Rio Grande above Hatch, valleys maintain their depth. Except near the apex, extensive surfaces have smooth desert pavement. On short, steep fans, gravel shows minimal weathering and are weakly cemented with caliche; age probably Wisconsinan and Holocene. On broad, more gently sloping fans, gravels are more weathered and commonly cemented by caliche; age probably pre-Wisconsinan. In south half of the state, gravel facies is characterized by creosote bush cover. Thin alluvial gravel covering pediments is denoted by Ig over subscript that identifies parent formation



Sand Facies — Sandy alluvium with subordinate amounts of fine gravel, silt, and clay. Forms at least four kinds of ground: 1) On short, steep fans sloping from the mountains of granitic or gneissic rock (e.g., parts of the Florida Mountains), this facies may form a smooth sandy layer a few feet thick covering gravel below; slopes 5 to 20 percent. 2) On other short fans, sand may form arcuate belt at toe of fan with slopes averaging 10 percent, commonly reworked into coppice dunes 3 to 7 ft high (sm). 3) Other belts of smooth sandy ground commonly slope 5 percent or less and consist of sand mounds approximately 1 ft high over caliche (ss₁). 4) Gypsiferous sand (ss₂), especially in the Jornada del Muerto, Tularosa Valley and east side of the Pecos Valley. Sand facies absent on the broad Las Palomas surface. Thin fan sand covering pediments is denoted by Is over subscript that identifies underlying formation. Boundary with residual sand, fan gravel, and fan silt is approximate



Silt Facies — In Basin and Range parts of the state, toes of fans may be silty and clayey rather than sandy; surface smooth, with slopes less than 5 percent. Slow infiltration rates and low slopes result in sluggish runoff. Forms a belt below the sand facies and grades downslope to plays silt (psl) with slopes less than 2 percent. Abundant swelling clays and exchangeable sodium. Surface layers predominantly Holocene; subject to sheet flooding, gradational with al₁. East and west of Sangre de Cristo Mountains, also forms fans of sandy or silty loam with little gravel in upper 3 to 4 ft, but abundant gravel below the loam. Caliche soft. Includes loess on isolated hillsides. Boundary with residual loam (rl), plays silt (psl), and fan sand (Is) approximate

EOLIAN DEPOSITS

Eolian deposits are laid down by wind, mostly as sheets of sand or silt (loess). Rarely, after prolonged drought on shale desert in the San Juan Basin, shale flags may accumulate in rippled sheets or even small dunes, but with the next rain, these become mud. Sand dune shapes depend on topography, relative strength of the winds, supply of sand, and vegetation. Some dunes are concave towards the windward (parabolic), others are concave towards the leeward (barhans), and others are longitudinal or transverse. Some dune clusters (e.g., Great White Sands) have all four kinds. Dunes may climb a windward slope or fill on a leeward slope. Most of New Mexico's eolian sand sheets have a basal layer of weathered, partly cemented, reddish stabilized sand, some sand surfaces on such layers are smooth. In the Basin and Range and Great Plains parts of the state, these surfaces are generally underlain by caliche; in the San Juan Basin, sand sheets commonly overlie residuum, fan deposits, or bedrock. Where sand is thick, as on sand facies of fans in the Basin and Range and at climbing dunes east of the Pecos River (Mescalero Sands) the sand is in mounds (coppice dunes) with profuse growth of vegetation — mesquite, and saltbush in the Basin and Range; sand sage, shinnery oak, small soapweed yucca, and occasional mesquite on the Mescalero Sands. Sand sheets are predominantly late Pleistocene; mounds and dunes are largely Holocene

s/b SAND UNDERLAIN BY BASALT — Extensive on basaltic plains south and east of Zuni Mountains and on West Potrillo Mountains. At Kilbourne Hole and Hunt's Hole, the sand is of volcanic origin

s/ca/OTs SAND UNDERLAIN BY CALICHE ON SANTA FE GROUP — Mostly on La Mesa and south part of the Jornada del Muerto

s₁/ca/To THIN SAND ON CALICHE ON OGALLALA FORMATION — Thickness about 1 ft. Chips of caliche comprise 30 percent of the sand. Generally too shallow for farming, but good shallow source for aggregates

s₂/ca/To MODERATELY THICK SAND ON CALICHE ON OGALLALA FORMATION — Sand 1 to 3 ft thick. Surface layers noncalcareous over reddish loam. Local sand mounds. Ground favorable for farming. Boundaries approximate

s₃/ca/To THICK SAND ON CALICHE ON OGALLALA FORMATION — Sand 3 to 5 ft thick. Local mounds. Brownish-red, fine sandy loam over reddish-brown, sandy clay loam; noncalcareous to depths of 3 ft; calcareous subsoil contains filaments of lime carbuncles. Where farmed, ground is subject to wind erosion. Boundaries approximate

sm LOOSE SAND IN MOUNDS — Coppice dunes, commonly 3 to 7 ft high and 25 to 50 ft in diameter; generally elongated north of east but a local exception lies east of Columbus where elongation is south of east. Age is Holocene. Boundaries fairly accurate

ss, s SAND SHEETS — Surfaces smooth except for ripples 2 to 3 inches high and scattered sand mounds 3 to 12 inches high, especially around small shrubs. Thickness of loose sand generally more than about 12 to 24 inches, but commonly overlies stabilized sand. Underlying material where known identified by subscript

ld LONGITUDINAL DUNES — Sand commonly 6 ft thick, locally 10 ft. Forms distinct ridges generally oriented north of east. Locations diagrammatic and width exaggerated

ds OTHER DUNES — ds₁, quartzose sand, ds₂, gypsiferous sand

l/b LOAM ON OLD BASALTIC LAVA — Probably pre-Wisconsinan loess

si EOLIAN SILT

EXPLANATION OF SURFICIAL GEOLOGY

by Charles B. Hunt 1977

New Mexico has two kinds of lake deposits in addition to those forming today in arid or semiarid areas. The most extensive deposits were laid down in Pleistocene lakes that flooded closed basins now marked by playas. Many of these deposits in the Basin and Range are of late Tertiary. Most numerous are the so-called "badland" deposits of the Great Plains and the Ogallala Formation. Some of these valleys and badland valleys with sand mounds on the lee side others may be due to salt in and up of the surface. Still others may be attributed to warping. Third are sinkholes clearly due to solution, like Bottomless Lake, sinks at Santa Rita and some of the depressions related to karst of the San Andres Formation and a low-lying ground north of the Sacramento Mountains. A fourth type is represented by ephemeral ponds in swales marking cutoff meanders on alluvial floodplains. A fifth type occurs only in the main valleys at Robinson Hole, Hunt's Hole, and Zuni Salt Lake. Only the first three types appear on the map. Area of deposits represented has been exaggerated because of map scale, but total area probably about right because smaller deposits are omitted.

PSI SILEY LAKE OR PLAYA DEPOSITS -- Ground mostly bare, gypsiferous deposits labeled ps₂.

PS SANDY LAKE OR PLAYA DEPOSITS -- Gypsiferous deposits labeled ps₂.

bw, dg, sp BEACH DEPOSITS -- Sand or gravel, sandy stretches mostly reworked into low dunes. Incompletely shown.

ev EVAPORITES -- Saline or alkaline deposits precipitated from brines in playas having high evaporation rates, notably Estancia Valley, Animas Valley, and Zuni Salt Lake. Salts are gradational with playa silt (psi) and occur in orderly concentric zones reflecting relative solubility of the salts. Thicknesses range from 1 to several inches, but salts mixed with mud may be tens of feet deep. Efflorescent crusts subject to wind erosion contribute to salinity of ground to leeward.

GLACIAL AND PERIGLACIAL DEPOSITS

During the Pleistocene New Mexico had mountain (alpine) glaciers high on the Sangre de Cristo Range, Tuzigoot Mountains, and Sierra Blanca Peak. The source of such glaciers was in nearly circular, steep-sided basins (cirques) at valley heads. High valleys eroded by the glacial tongues tend to be U-shaped; at lower elevations where eroded by streams, these valleys are V-shaped. Gravels deposited along each side of valley ice represent debris that rolled down the mountainside onto the ice to form lateral moraines. Hummocky ridges of sand and gravel deposited across the lower ends of the glaciers form terminal moraines. Within the cirques generally stand two ramparts of boulders. An inner rampart, forming only, is located at the lower edge of the snowfield that accumulated annually in the cirque. It represents rocks broken by frost from the headwall of the cirque, rolled down the snowbank, and collected at the ridge. Their inner ridges are peders. Farther out in the cirque -- perhaps at the mouth -- is a second ridge, covered with fine unweathered rock darkly stained with iron and manganese oxides. These outer ridges probably formed during the mid-Pleistocene "little ice age".

mg MOUNTAIN GLACIERS -- Extent exaggerated.

pg PERIGLACIAL DEPOSITS OR MOUNTAIN TOPS -- Primarily represented by boulder fields and patterned ground where frost action was intensive during the glaciations. Extent and boundaries approximate; graded laterally to stony retiduum and colluvium.

av AVALANCHE DEPOSITS -- Bouldery; some are lag concentrates of boulders where fine grained sediments have been removed by erosion. Deposits narrow and long downslope; commonly 10 to 50 ft thick. Apparently deposited as mudflows during late Pleistocene time when there were numerous perennial mountain snowfields. Frost action at the time was vigorous; sudden thaws could trigger floods or mudflows on the mountainsides. Slow movement downslope may be reactivated in artificial cuts through these deposits if water enters the plane of slippage.

lds LANDSLIDE DEPOSITS -- Abundant on slopes of Cretaceous shale. Whereas avalanche deposits are elongate downslope, landslide deposits are short downslope but wide along the contour. Characteristically, they retain a cap of the lava or sandstone sloping into the hillside atop a steep colluvial covered shale slope. Stabilized landslides may be reactivated if water is allowed to enter the plane of slippage.

MISCELLANEOUS TYPES OF GROUND

ba BASALT -- Includes lava flows, lava cones, cones of scoriae, necks, and fields of scoriae. Presumably Quaternary and late Tertiary; some young enough to have retained original weathering and retained their original structures and shapes. are commonly referred to as malpais (Spanish, bad ground). Includes some Tertiary basalt that conspicuously controls the topography. Locally covered by loam (lb), eolian deposits, silt, stream deposits). These older surfaces are more deeply eroded, tilted, and faulted. Individual flows generally less than 50 ft thick; locally, several flows may aggregate a few hundred feet thick. Commonly interbedded with volcanic ash (tuff). Excludes lavas mantled by loess or other sediments; such areas indicated by subscript (e.g., lbh -- loam over basalt; fsb -- fan sand over basalt). Boundaries shown are adequate.

o OTHER BEDROCK -- Colluvium or other cover amounts to less than half the area. Only extensive areas are shown; age and rock type keyed by symbol to State geologic map (e.g., Kd, Cretaceous Dakota Sandstone; Rk Triassic Santa Rosa Sandstone). Many small areas omitted; indicated boundaries are approximate. Principal formations and subscripts used are:

Og -- Gatuna Fm.
Oht -- Bandelier Tuff
Ovt -- Rhyolite Flows
Q1-C -- Upper Santa Fe Group
Q1-S -- Santa Fe Group, undivided, and related formations
Q1g -- Gila Conglomerate
Tc -- Ogallala Fm.
Tc1 -- Lower Santa Fe Group
Tc2 -- Chuska Sandstone
Tc3 -- Altiplano and lacustrine deposits
Tc4 -- Cretaceous conglomerate (generally equivalent to Las Pintas Fm.)
Tpi -- Pecos Tuff
Tp -- Pecos volcanic series
T5 -- Tertiary volcanics, largely Datil Fm. in SW, includes some pre- and post-Datil volcanic sequences
Tsh -- Blanca Mesa Fm.
Tg -- Galisteo Fm.
Tj -- San Jose Fm.
Tn -- Mesquite Fm.
T -- Tertiary volcanics, largely Datil Fm. in SW, includes some pre- and post-Datil volcanic sequences
Tpc -- Pecos Canyon Fm.
Tka -- Animas Fm.
TKt -- Raton Fm.
TKs -- Rio Alamo Sandstone
Kv -- Volcanics of Cretaceous age; various composition
Kkf -- Kirtland Shale and Fruitland Fm.
Kpc -- Pictured Cliffs Sandstone
Kl -- Lewis Shale
Kmv -- Cretaceous sandstone and shale, mostly Mesaverde Fm.
Ksh -- Cliffhouse Sandstone
Kpl -- Point Lookout Sandstone
Ksh -- Cretaceous shale
Kg -- Gallup Sandstone
Km -- Mancos Shale
Kd -- Dakota Sandstone
J -- Jurassic, undivided
Jm -- Morrison Fm.
Jz -- Zuni Sandstone
R, J -- Triassic and Jurassic, undifferentiated
R -- Triassic, undifferentiated
Rgc -- Glen Canyon Sandstone
R -- Chinle Fm.
R -- Santa Rosa Sandstone
R -- Galisteo Fm.
R -- Animas Group
R -- San Andres Fm. (limestone)
R -- Dakota Sandstone
R -- Permian Fm.

Py -- Yozo Fm.
Pa -- Abo Fm.
Ph -- Hueco Fm.
Pal -- Paleozoic, undivided
Pms -- Madera Limestone and Sandia Fm., undivided
P, P' -- Permian, Pennsylvanian
M, D -- Mississippian, Devonian
S, O, E -- Silurian, Ordovician, Cambrian
pC -- Precambrian
gr -- Granitic, gneissic, and intrusive rocks of various ages

D Disturbed ground. Mostly urban areas large enough to show on state base; farmed lands excluded. Includes airports, mined areas, tailings dumps, and feedlots. Incompletely shown.

X Open pits for road fill, sand, gravel, caliche, or other aggregates.

o Playa-lake depressions. Mostly small closed basins produced by eolian activity and local solution subsidence.

REFERENCES

Dane, C.H., and Bachman, G.O., 1965, Geologic map of New Mexico: U.S. Geological Survey, Washington, D.C.

Hawley, J.W., Bachman, G.O., and Manley, Kim, 1976, Quaternary stratigraphy in the Basin and Range, and Great Plains provinces, New Mexico and Western Texas, in The Quaternary stratigraphy of North America, W.C. Mahaney, ed: Stroudsburg, Pennsylvania, Dowden, Hutchinson and Ross, p. 235-274.

New Mexico State University, Agricultural Experiment Station, Research reports showing soil association and land classification for irrigation for each county.


New Mexico State Highway Department supplied data for aggregate resources in New Mexico.

Soil Conservation Service, 1/62,500 aerial mosaics of New Mexico Quadrangles.


Data from these and other sources were plotted on the 1/250,000 quadrangle maps, field checked with about 40,000 mi of automobile traverses and 20 hours aerial reconnaissance over areas difficult of ground access. Mapping began spring 1974 and was completed June 1976.

ACKNOWLEDGMENTS

The author wishes to thank John W. Hawley and Robert H. Weber of the New Mexico Bureau of Mines and Mineral Resources for critically reviewing the maps and explanation; also Neils M. Pearson, for editing the explanation and for handling total cartographic compilation.



Index map of New Mexico



YUCCA PLANTS

INTRODUCTION

Surficial geology concerns origin, distribution, and significance of deposits and soils at or near the surface. Completely bare bedrock forms probably less than 5 percent of New Mexico's land surface; consequently surficial materials form by far the largest and most-used part of the ground around us. Several aspects of surficial geology that contribute significantly to an understanding of our environment are water yielding properties of the ground; its susceptibility to flooding and erosion; its susceptibility to such hazards as landslides, avalanches, and earthquakes; ease of excavation; suitability for foundations and road building; agricultural potential, including suitability for irrigation or pasturage; and mineral resources potential.

Surficial materials commonly are poorly consolidated, consisting partly of bedrock weathered in situ (residuum), but mostly of sediments derived by erosion and transported by water, wind, ice, or gravity (mass wasting) to a site of temporary deposition before being further eroded and transported downslope.

Four major categories of surficial materials are distinguished on the map by color: residual materials, transitional deposits, transported deposits, and miscellaneous types of ground.

RESIDUAL MATERIALS

Materials generally formed in place, including: residuum, formed in situ by weathering of a parent formation; caliche; travertine and related spring deposits; shale or sandstone baked by coal beds burning in situ (clinker); karst and related deposits in sinks, and the following, which are not distinguished on the map -- organic deposits; desert pavement; cave deposits; and desert varnish.

RESIDUUM

In New Mexico, residuum tends to be thin, generally less than 2 ft thick -- rarely as much as 5 ft. Texture depends upon composition of parent rock, and ranges from clay to coarse sand; texture may be bouldery in granitic areas. Areas shown as residuum include small outcrops of parent rocks and some alluvial or eolian deposits either mistaken for residuum or too small to show on the map. These materials are predominantly of late Pleistocene (Wisconsinan) or Holocene age. Ground is hummocky with slopes less than 10 percent; scattered small outcrops of resistant beds form small ledges.

LOAMY RESIDUUM — Texture variable -- mixed clay, silt, and sand. Thickness 1 to 5 ft. Parent formations fine grained, shallow, and identified by subscripts. Where clayey, this residuum generally contains appreciable amounts of swelling clay and is highly susceptible to sodium exchange, especially over the Chinle Formation (subscript Tlc), Cretaceous shale (subscript Ksh), and Tertiary clayey volcanic formations. Slopes locally 10 percent and subject to washing. Although the unit is distinctive, the indicated boundaries are approximate.

STONY RESIDUUM — Stony residuum, with accompanying sand and silt. Thickness mostly less than 3 ft. Texture variable depending on parent material, indicated by subscript. Boundaries gradational with cl and lg.

STONY LOAM OVER BASALT — Lithology highly variable; locally abundant clay and silt, probably local; stones basaltic, mostly rough scoriae or angular blocks and flakes. Includes alluvium along small washes; numerous basalt mounds and low scarps along some washes and at edges of flows; thickness generally less than 3 ft. Surface smooth; slopes usually less than 5 percent except at sides of washes, bases of volcanic cones (including spatter cones), and edges of flows. Not subject to severe erosion. Boundaries indicated are fairly well defined despite variable lithology; boundaries with alluvium are approximate.

SANDY OR SANDY LOAM RESIDUUM — The shallow sandy or sandy silt substrates are distinguished by subscripts (e.g., rs/Kd, sandy residuum over Dakota Sandstone). Thickness commonly 1 ft. Subject to wind erosion where vegetation is sparse; minimal washing. A distinctive unit with adequate boundaries, except in the San Juan Basin and along the Canadian River.

GYPSEOUS AND SANDY RESIDUUM ALONG PECOS RIVER VALLEY — Parent material Artesia (Pal) and related formations. Rarely over 2 ft thick. Numerous small outcrops of gypsum thinly mantled by loose sand with or without small pebbles. A distinctive unit; boundaries are approximate.

RESIDUUM ON LIMESTONE — Widespread on east slope of Sacramento Mountains, Chupadera Mesa, and flanks of Zuni Mountains; less extensive on Cretaceous limestone beds south of Raton. Stony and blocky; generally well cemented with calcium carbonate; little subject to erosion. Slopes average steeper than most residuum. Thickness generally less than 2 ft, rarely as much as 5 ft. A distinctive unit; boundaries indicated are adequate.

CALICHE

CALICHE — Partly indurated zone of calcium carbonate accumulation formed in upper layers of surficial deposits; 2 to 10 ft thick; commonly overlain by windblown sand. Much caliche shown on the map consists of tough, slabby surface layers underlain by calcium carbonate nodules that grade downward to fibers and veinlets. Especially well developed in Basin and Range and Great Plains parts of the state. Thick caliches (locally >20 ft) associated with undissected High Plains surfaces of the Great Plains commonly comprise an upper sequence of several carbonate-cemented zones interlayered with reddish loamy caliche horizons over a basal caprock zone developed on Ogallala (Tn) sediments. Forms on various types of parent formations, indicated by subscripts. The extensive caliche along Rio Salado northwest of Socorro is partly a travertine deposit. Where buried by sand, the caliche is identified by subscript ca. A distinctive unit; boundaries are well defined where the caliche forms rimrock and approximate where exposed in deflation hollows. Where thick and well indurated, caliche is quarried for road metal and other aggregate, subject to minimal erosion.

SPRING DEPOSITS

TRAVERTINE AND RELATED DEPOSITS — Most deposits shown have been formed at springs discharging water hotter than 100°F (34°C). Travertine mounds and benches to 50 ft high. Deposits at east base of Mesa Lucero may not have been created by hot springs.

CLINKER

SLAGGY COAL ASH AND VITRIFIED SHALE AND SANDSTONE MASSES FUSED BY BURNING COAL BEDS — Incompletely shown -- coal may ignite spontaneously, by lightning or ground fire. Depending on oxygen availability, the coal may burn tens of feet back into the ground. Common in coal-bearing formations of San Juan Basin and Raton district. Used for road metal.

KARST DEPRESSION DEPOSITS

KARST-RELATED DEPOSITS — Underground solution of limestone and gypsum produces caverns or smaller subsurface voids, and causes roof-rock collapse, forming closed karst depressions (sinkholes) at the surface, mantled with blocks of the roof rock. Widespread in San Andres Formation (subscript Pca) north of the Sacramento Mountains and on Chupadera Mesa. Sinks commonly 50 ft deep and 500 to 1,000 ft wide. Similar deposits composed of slumped gravel and alluvium along the Pecos River valley are attributed to solution of underlying gypsum or other salts. Slumped beds dip 1 to 5 degrees into the depression; may be overlain by undisturbed gravels. Thickness to 300 ft. Although these are distinctive features, extent and boundaries, largely derived from the 1/250,000 quadrangle maps, are approximate.